

Student Name: Jane R.

Pt Initials: S.L.

Pt Allergies: PCN, Percocet, ASA, tape

Medication List

Drug (Generic, Trade and Class)	Dosage/Route/ Frequency	Rationale	Nursing Implications
Diltiazem (Cardizem): CCB	60 mg po BID	Angina	Assess location, duration, intensity & precipitating factors of anginal pain
Metoprolol (Lopressor): antianginal, antihypertensive (beta blocker)	100mg po QD	Post MI	Take apical pulse before administering. If <50bpm or arrhythmia, hold mx & notify MD; administer with meals or p eating. ER tabs swallow whole, do not cursh, break or chew
Metronidazole (Flagyl): anti-infective, anti-protozoal, antiulcer	250mg po Q8hrs X7days	Osteomyelitis of right foot	Administer with food or milk to minimize GI irritation; may crush tablets
Regular SSI	SSI AC & HS	Pt receiving high dose steroids	Onset: 30-60min Peak: 2-4hr Duration: 5-7hr Assess pt for s/sx of hypoglycemia Do not mix insulin glargine with any other insulin
Furosemide(Lasix): loop diuretic	40 mg IV BID	Pulmonary edema	Administer undiluted. Rate: slowly over 1-2 min Administer in AM Monitor I&O